

BIOLOGICALLY ACTIVE 4H-BENZO[1,4]OXAZIN-3-ONES

Cross Reference to Related Applications

5 This application is a continuation-in-part of U.S. Serial No. 09/854,302, ^{now U.S. Patent No. 6,555,536} filed May 11, 2001, which in turn claims priority from U.S. Serial No. 60/203,860, filed May 12, 2000.

10 Field of the Invention

This invention relates to novel 4H-benzo[1,4]oxazin-3-ones useful for the treatment of Non-Insulin Dependant Diabetes Mellitus (NIDDM) and complications thereof and disorders related to lipid metabolism and energy homeostasis such as obesity. More particularly, the compounds act through the peroxisome proliferator activated receptor gamma (PPAR γ). Compounds in the series are PPAR γ modulators.

20 Background of the Invention

Diabetes is a disease caused by multiple factors and characterized by hyperglycemia which may be associated with increased and premature mortality due to an increased risk for microvascular and macrovascular diseases such as nephropathy, neuropathy, retinopathy, atherosclerosis, polycystic ovary syndrome (PCOS), hypertension, ischemia, stroke, and heart disease. Type I diabetes (IDDM) results from genetic deficiency of insulin, the hormone regulating glucose metabolism. Type II diabetes is known as non-insulin dependent diabetes mellitus (NIDDM), and is due to a profound resistance to insulin regulatory effect on glucose and lipid metabolism

in the main insulin-sensitive tissues, i.e., muscle, liver and adipose tissue. This insulin resistance or reduced insulin sensitivity results in insufficient insulin activation of glucose uptake, oxidation and storage in muscle and inadequate insulin repression of lipolysis in adipose tissue as well as glucose production and secretion in liver. Many Type II diabetics are also obese, and obesity is believed to cause and/or exacerbate many health and social problems such as coronary heart disease, stroke, obstructive sleep apnoea, gout, hyperlipidemia, osteoarthritis, reduced fertility, and impaired psychosocial function.

A class of compounds, thiazolidinediones (glitazones), have been suggested to be capable of ameliorating many symptoms of NIDDM by binding to the peroxisome proliferator activated receptor (PPAR) family of receptors. They increase insulin sensitivity in muscle, liver and adipose tissue in several animal models of NIDDM resulting in correction of the elevated plasma levels of glucose, triglycerides and nonesterified free fatty acids without any occurrence of hypoglycemia. However, undesirable effects have occurred in animal and/or human studies including cardiac hypertrophy, hemadilution and liver toxicity.

Most PPAR γ agonists currently in development have thiazolidinedione ring as their common chemical structure. PPAR γ agonists have been demonstrated to be extremely useful for the treatment of NIDDM and other disorders involving insulin resistance. Recently, troglitazone, rosiglitazone, and pioglitazone have been approved for treatment of type II diabetes. There is also indication that benzimidazole-containing

ORT-1527

thiazolidinedione derivatives may be used to treat irritable bowel disorder (IBD), inflammation, and cataract (JP 10195057).


- 5 JP 09012576 (Yoshitake et al.) discloses benzothiazine derivatives stated as useful therapeutic agents for circulatory system disease and glaucoma.

- 10 JP 09012575 (Hiroaki et al.) discloses benzoxazine and benzothiazine derivatives stated to be useful as prophylactic drugs and/or therapeutic drugs in hyperlipemia, hyperglycemia, obesity, diseases attributable to sugar tolerance insufficiency, hypertension, osteoporosis, cachexia, and complications
15 of diabetes such as retinopathy, nephrosis, neuropathy, cataract, coronary artery disease and arteriosclerosis.

- 20 WO 99/20614 (Lohray et al.) discloses β -aryl- α -oxysubstituted alkylcarboxylic acids stated as antiobesity and hypocholesterolemic compounds which may have agonist activity against PPAR α and/or PPAR γ , and optionally inhibit HMG CoA reductase.


- 25 WO 97/17333 (Frechette et al.) and U.S. Patent Nos. 5,696,117 and 5,854,242 to Frechette et al. disclose benzoxazine and pyrido-oxazine compounds having a moiety of a fused phenyl or fused pyridyl, pharmaceutical compositions containing the compounds, and methods for their production and their use in treating bacterial
30 infections.

U.S. Patent No. 5,859,051 to Adams et al. discloses the following acetylphenols,



wherein substituents are as described in the reference,
which are stated to be useful as antiobesity and
antidiabetic compounds without the thiazolidinedione
moiety.


WO 99/38845 (De La Brouse-Elwood et al.) discloses
the following compounds,



wherein substituents are as described in the reference,
which are stated to modulate the PPAR γ receptor and are
stated as useful in the diagnosis and treatment of type
II diabetes (and complications thereof) and inflammatory
disorders.

Summary of the Invention

The present invention is directed to a compound of
Formula I,



I

or an optical isomer, enantiomer, diastereomer, racemate
or racemic mixture thereof, ester, prodrug form, or a
pharmaceutically acceptable salt thereof, wherein

A is selected from aryl, heterocyclyl, and C₁-C₁₀ alkyl, said aryl, heterocyclyl, and C₁-C₁₀ alkyl being optionally substituted with one or more members selected from the group consisting of halogen, OH, aryl, C₃-C₈ cycloalkyl, C₁-C₁₀ alkyl substituted with a halogen, C₁-C₁₀ alkyl ether, heterocyclyl, carbonyl, oxime, -N(R¹)(SO₂R), -C(NNR³R⁴)R¹, -COOR¹, -CONR¹R², -OC(O)R¹, -OC(O)OR¹, -OC(O)NR¹R², -NR¹R², -NR³C(O)R¹, -NR³C(O)OR¹, and -NR³C(O)NR¹R², wherein

10 R is selected from C₁-C₆ alkyl, trifluoromethyl, phenyl, and substituted phenyl;

15 R¹ and R² are independently selected from hydrogen, C₁-C₁₀ alkyl, aryl, heterocyclyl, and alkylaryl, or R¹ and R² may be taken together to form a 5- to 10-member ring; and

20 R³ and R⁴ are independently selected from hydrogen, C₁-C₁₀ alkyl, aryl, heterocyclyl, alkylaryl, -C(O)R¹, or -C(O)NR¹R²;

25 Z¹ is selected from hydrogen, C₁-C₆ alkyl, aryl, heterocyclyl, COOR¹, CONR¹R², OH, C₁-C₆ alkyl ether, -OC(O)R¹, -OC(O)OR¹, -OC(O)NR¹R², -NR¹R², -NR³C(O)R¹, -NR³C(O)OR¹, -NR³C(O)NR¹R², halogen, -C(O)R¹, -C(NR³)R¹, -C(NOR³)R¹, and -C(NNR³R⁴)R¹;


Z² is selected from hydrogen, halogen, C₁-C₆ alkyl;

30 Z¹ and Z² may together form a fused aromatic ring;

n is an integer from 0 to 3;


ORT-1527

G is selected from $-\text{COOR}^1$, $-\text{C}(\text{O})\text{COOR}^1$, $-\text{CONR}^1\text{R}^2$, $-\text{CF}_3$, $-\text{P}(\text{O})(\text{OR}^1)(\text{OR}^2)$, $-\text{S}-\text{R}^8$, $-\text{O}-\text{R}^8$,




R^5 and R^6 are independently hydrogen or $\text{C}_1\text{-C}_6$ alkyl;
 R^7 is hydrogen, $\text{C}_1\text{-C}_6$ alkyl, or $-\text{C}(\text{O})\text{R}^5$;
 R^8 is selected from the group consisting of hydrogen, $\text{C}_1\text{-C}_6$ alkyl, and substituted $\text{C}_1\text{-C}_6$ alkyl; and
 B is oxygen or $-\text{NR}^5$;

E is selected from hydrogen, $\text{C}_1\text{-C}_6$ alkyl and a moiety of the formula




X is hydrogen or oxygen, with the proviso that

when E is hydrogen and G is $-\text{COOH}$, $-\text{COOCH}_3$, or a moiety of the formula of



A is selected from the group consisting of aryl, heterocyclyl, substituted C₁-C₆ alkyl and C₇-C₁₀ alkyl, provided that when X is hydrogen, n is 1 and G is a moiety of the formula of



5

A is selected from the group consisting of heterocyclyl, and C₇-C₁₀ alkyl.

The compounds of the present invention are PPAR γ modulators useful for the treatment of NIDDM and complications thereof and disorders related to lipid metabolism and energy homeostasis such as obesity.

Illustrative of the invention is a pharmaceutical composition comprising a pharmaceutically acceptable carrier and any of the compounds described above. Illustrating the invention is a pharmaceutical composition made by mixing any of the compounds described above and a pharmaceutically acceptable carrier. An illustration of the invention is a process for making a pharmaceutical composition comprising mixing any of the compounds described above and a pharmaceutically acceptable carrier.

An embodiment of the invention is a method of treating a subject suffering from a disorder in glucose and lipid metabolism, which comprises administering to the subject a therapeutically effective amount of a compound of Formula I.

30

Another embodiment of the invention is a method of inhibiting the onset of a condition of a disorder in glucose and lipid metabolism, which comprises administering to the subject a prophylactically effective
 5 dose of a compound of Formula I.


Further illustrating the invention is a method of treating a subject suffering from a disorder in glucose and lipid metabolism, which comprises administering to
 10 said subject an effective amount of a compound of Formula I, or an optical isomer, enantiomer, diastereomer, racemate or racemic mixture thereof, ester, prodrug form, or a pharmaceutically acceptable salt thereof,, wherein said disorder is selected from NIDDM, obesity,
 15 nephropathy, neuropathy, retinopathy, atherosclerosis polycystic ovary syndrome, hypertension, ischemia, stroke, heart disease, irritable bowel disorder, inflammation, and cataract.

Also included in the invention is the use of any of the compounds described above for the preparation of a medicament for treating a condition selected from NIDDM, obesity, nephropathy, neuropathy, retinopathy, atherosclerosis polycystic ovary syndrome, hypertension,
 20 ischemia, stroke, heart disease, irritable bowel
 25 disorder, inflammation, and cataract in a subject in need thereof.

Detailed Description of the Invention

30

The present invention provides 4H-benzo[1,4]oxazin-3-ones of Formula I,



I

or an optical isomer, enantiomer, diastereomer, racemate or racemic mixture thereof, ester, prodrug form, or a pharmaceutically acceptable salt thereof, wherein

5

A is selected from aryl, heterocyclyl, and C₁-C₁₀ alkyl, said aryl, heterocyclyl, and C₁-C₁₀ alkyl being optionally substituted with one or more members selected from the group consisting of halogen, OH, aryl, C₃-C₈ cycloalkyl, C₁-C₁₀ alkyl substituted with a halogen, C₁-C₁₀ alkyl ether, heterocyclyl, carbonyl, oxime, -N(R¹)(SO₂R), -C(NNR³R⁴)R¹, -COOR¹, -CONR¹R², -OC(O)R¹, -OC(O)OR¹, -OC(O)NR¹R², -NR¹R², -NR³C(O)R¹, -NR³C(O)OR¹, and -NR³C(O)NR¹R², wherein

10

15

R is selected from C₁-C₆ alkyl, trifluoromethyl, phenyl, and substituted phenyl;

R¹ and R² are independently selected from hydrogen, C₁-C₁₀ alkyl, aryl, heterocyclyl, and alkylaryl, or R¹ and R² may be taken together to form a 5- to 10-member ring; and

20

R³ and R⁴ are independently selected from hydrogen, C₁-C₁₀ alkyl, aryl, heterocyclyl, alkylaryl, -C(O)R¹, or -C(O)NR¹R²;

25

Z¹ is selected from hydrogen, C₁-C₆ alkyl, aryl, heterocyclyl, COOR¹, CONR¹R², OH, C₁-C₆ alkyl ether, -OC(O)R¹, -OC(O)OR¹, -OC(O)NR¹R², -NR¹R², -NR³C(O)R¹, -NR³C(O)OR¹, -NR³C(O)NR¹R², halogen, -C(O)R¹, -C(NR³)R¹, -C(NOR³)R¹, and -C(NNR³R⁴)R¹;

30

[illegible]

```
5  n is an integer from 0 to 3;
```

```
5  n is an integer from 0 to 3;
```

G is selected from $-\text{COOR}^1$, $-\text{C}(\text{O})\text{COOR}^1$, $-\text{CONR}^1\text{R}^2$, $-\text{CF}_3$, $-\text{P}(\text{O})(\text{OR}^1)(\text{OR}^2)$, $-\text{S}-\text{R}^8$, $-\text{O}-\text{R}^8$,



R⁷ is hydrogen, C₁-C₆ alkyl, or -C(O)R⁵;


15

R is selected from hydrogen, C₁-C₆ alkyl and a moiety of the formula




10

when E is hydrogen and G is -COOH, -COOCH₃, or a moiety of the formula of




A is selected from the group consisting of aryl, heterocyclyl, substituted C₁-C₆ alkyl and C₇-C₁₀ alkyl, provided that when X is hydrogen, n is 1 and G is a moiety of the formula of



A is selected from the group consisting of heterocyclyl, and C₇-C₁₀ alkyl.

Preferably, compounds of Formula I, wherein X is oxygen are embodiments of the present invention.


Compounds of Formula I wherein E is C₁-C₆ alkyl or a moiety of the formula




are also preferred embodiments of this invention.

Compounds of Formula I wherein


A is optionally substituted C₁-C₆ alkyl or optionally substituted aryl;
X is oxygen; and

G is selected from $-\text{COOR}^1$, $-\text{CONR}^1\text{R}^2$, $-\text{CF}_3$, , -

$\text{P}(\text{O})(\text{OR}^1)(\text{OR}^2)$, $-\text{S}-\text{R}^8$, $-\text{O}-\text{R}^8$, and 

are preferred embodiments of the present invention as well. Particularly, A is substituted C_1 - C_6 alkyl and G is COOH or COOCH_3 .

The following compounds are still preferred embodiments of the present invention:



Unless otherwise noted, "alkyl" and "alkoxy" as used herein, whether used alone or as part of a substituent group, include straight and branched chains having 1 to 10 carbon atoms, or any number within this range, optionally substituted with one or more independent groups including, but not limited to, H, halogen (F, Cl,

Br, I), C₁-C₁₀ alkyl, OH, amino, alkoxy, alkthio, aryl, substituted aryl, heterocyclyl, substituted heterocyclyl, COOR¹, and CONR¹R² wherein R¹ and R² are as described above. For example, alkyl radicals include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, t-butyl, n-pentyl, 3-(2-methyl)butyl, 2-pentyl, 2-methylbutyl, neopentyl, n-hexyl, 2-hexyl, 2-methylpentyl, and the like. Alkoxy radicals are oxygen ethers formed from the previously described straight or branched chain alkyl groups. Cycloalkyl groups contain 3 to 8 ring carbons and preferably 5 to 7 ring carbons. Similarly, alkenyl and alkynyl groups include straight and branched chain alkenes and alkynes having 1 to 10 carbon atoms, or any number within this range.

The term "Ar" represents aryl. Unless otherwise stated, "aryl," employed alone or in combination with other terms (e.g., aryloxy, arylthioxy, arylalkyl), is an aromatic radical which can be a single ring or multiple rings which are fused together or linked covalently. Illustrative aryl groups may be phenyl or naphthyl optionally substituted with one or more independent groups such as H, halogen, C₁-C₁₀ alkyl, C₃-C₈ cycloalkyl, COOR¹, CONR¹R², OH, C₁-C₁₀ alkyl ether, aryl or heterocyclyl ether, OC(O)R¹, OC(O)OR¹, OC(O)NR¹R², NR¹R², NR³C(O)R¹, NR³C(O)OR¹, and NR³C(O)NR¹R² wherein R¹ and R² are as described above. When Z¹ and Z² as described above together form a fused aryl ring, the aromatic ring is fused to the phenyl moiety of benzoxazin in Formula I.

"Heterocyclyl" or "heterocycle" is a 3- to 8-member saturated, partially saturated, or unsaturated single or fused ring system which consists of carbon atoms and from one to three heteroatoms selected from N, O and S. The

heterocyclyl group may be attached at any heteroatom or carbon atom which results in the creation of a stable structure. Examples of heterocyclyl groups include, but are not limited to pyridine, pyrimidine, oxazoline, pyrrole, imidazole, morpholine, furan, indole, benzofuran, pyrazole, pyrrolidine, piperidine, and benzimidazole. "Heterocyclyl" or "heterocycle" may be substituted with one or more independent groups including, but not limited to, H, halogen, oxo, OH, C₁-C₁₀ alkyl, amino, and alkoxy.

It is intended that the definition of any substituent or variable at a particular location in a molecule be independent of its definitions elsewhere in that molecule. It is understood that substituents and substitution patterns on the compounds of this invention can be selected by one of ordinary skill in the art to provide compounds that are chemically stable and that can be readily synthesized by techniques known in the art as well as those methods set forth herein.

As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combinations of the specified ingredients in the specified amounts.

The term "subject" as used herein, refers to an animal, preferably a mammal, most preferably a human, who has been the object of treatment, observation or experiment.

Methods are known in the art for determining therapeutically and prophylactically effective doses for

the instant pharmaceutical composition. The term "therapeutically effective amount" as used herein, means that amount of active compound or pharmaceutical agent that elicits the biological or medicinal response in a tissue system, animal or human that is being sought by a researcher, veterinarian, medical doctor or other clinician, which includes alleviation of the symptoms of the disease or disorder being treated.

The term "prophylactically effective amount" refers to that amount of active compound or pharmaceutical agent that inhibits in a subject the onset of a disorder as being sought by a researcher, veterinarian, medical doctor or other clinician, the delaying of which disorder is mediated by the modulation of PPAR γ activity.

Depending upon the biological environment (e.g., cell type, pathological condition of the host, etc.), these compounds can activate or block the actions of PPAR γ . The utility of the compounds to treat disorders in glucose and lipid metabolism can be determined according to the procedures described herein. The present invention therefore provides a method of treating disorders in glucose and lipid metabolism in a subject in need thereof which comprises administering any of the compounds as defined herein in a quantity effective to such disorders. The compound may be administered to a patient by any conventional route of administration, including, but not limited to, intravenous, oral, subcutaneous, intramuscular, intradermal and parenteral.

The present invention also provides pharmaceutical compositions comprising one or more compounds of this

- invention in association with a pharmaceutically acceptable carrier. The pharmaceutical composition may contain between about 0.1 mg and 1000 mg, preferably about 100 to 500 mg, of the compound, and may be constituted
- 5 into any form suitable for the mode of administration selected. Carriers include necessary and inert pharmaceutical excipients, including, but not limited to, binders, suspending agents, lubricants, flavorants, sweeteners, preservatives, dyes, and coatings.
- 10 Compositions suitable for oral administration include solid forms, such as pills, tablets, caplets, capsules (each including immediate release, timed release and sustained release formulations), granules, and powders, and liquid forms, such as solutions, syrups, elixirs,
- 15 emulsions, and suspensions. Forms useful for parenteral administration include sterile solutions, emulsions and suspensions.

- To prepare the pharmaceutical compositions of this
- 20 invention, one or more compounds of Formula I or salt thereof of the invention as the active ingredient, is intimately admixed with a pharmaceutical carrier according to conventional pharmaceutical compounding techniques, which carrier may take a wide variety of
- 25 forms depending on the form of preparation desired for administration, e.g., oral or parenteral such as intramuscular. In preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed. Thus, for liquid oral preparations, such as
- 30 for example, suspensions, elixirs and solutions, suitable carriers and additives include water, glycols, oils, alcohols, flavoring agents, preservatives, coloring agents and the like; for solid oral preparations such as, for example, powders, capsules, caplets, gelcaps and

tablets, suitable carriers and additives include starches, sugars, diluents, granulating agents, lubricants, binders, disintegrating agents and the like. Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are obviously employed. If desired, tablets may be sugar coated or enteric coated by standard techniques. For parenterals, the carrier will usually comprise sterile water, though other ingredients, for example, for purposes such as aiding solubility or for preservation, may be included. Injectable suspensions may also be prepared, in which case appropriate liquid carriers, suspending agents and the like may be employed. The pharmaceutical compositions herein will contain, per dosage unit, e.g., tablet, capsule, powder, injection, teaspoonful and the like, an amount of the active ingredient necessary to deliver an effective dose as described above. The pharmaceutical compositions herein will contain, per unit dosage unit, e.g., tablet, capsule, powder, injection, suppository, teaspoonful and the like, from about 0.01 mg to 30 mg/kg of body weight per day. Preferably, the range is from about 0.03 to about 15 mg/kg of body weight per day, most preferably, from about 0.05 to about 10 mg/kg of body weight per day. The compounds may be administered on a regimen of 1 to 2 times per day. The dosages, however, may be varied depending upon the requirement of the patients, the severity of the condition being treated and the compound being employed. The use of either daily administration or post-periodic dosing may be employed.

Preferably these compositions are in unit dosage forms such as tablets, pills, capsules, powders, granules,

sterile parenteral solutions or suspensions, metered aerosol or liquid sprays, drops, ampoules, auto-injector devices or suppositories; for oral parenteral, intranasal, sublingual or rectal administration, or for administration by inhalation or insufflation. Alternatively, the composition may be presented in a form suitable for once-weekly or once-monthly administration; for example, an insoluble salt of the active compound, such as the decanoate salt, may be adapted to provide a depot preparation for intramuscular injection. For preparing solid compositions such as tablets, the principal active ingredient is mixed with a pharmaceutical carrier, e.g. conventional tableting ingredients such as corn starch, lactose, sucrose, sorbitol, talc, stearic acid, magnesium stearate, dicalcium phosphate or gums, and other pharmaceutical diluents, e.g. water, to form a solid preformulation composition containing a homogeneous mixture of a compound of the present invention, or a pharmaceutically acceptable salt thereof. When referring to these preformulation compositions as homogeneous, it is meant that the active ingredient is dispersed evenly throughout the composition so that the composition may be readily subdivided into equally effective dosage forms such as tablets, pills and capsules. This solid preformulation composition is then subdivided into unit dosage forms of the type described above containing from 0.1 to about 500 mg of the active ingredient of the present invention. The tablets or pills of the novel composition can be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged action. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope over the former. The two components can be separated by an enteric layer which

serves to resist disintegration in the stomach and permits the inner component to pass intact into the duodenum or to be delayed in release. A variety of material can be used for such enteric layers or coatings, such materials including a number of polymeric acids with such materials as shellac, cetyl alcohol and cellulose acetate.

The liquid forms in which the novel compositions of the present invention may be incorporated for administration orally or by injection include, aqueous solutions, suitably flavoured syrups, aqueous or oil suspensions, and flavoured emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil or peanut oil, as well as elixirs and similar pharmaceutical vehicles. Suitable dispersing or suspending agents for aqueous suspensions, include synthetic and natural gums such as tragacanth, acacia, alginate, dextran, sodium carboxymethylcellulose, methylcellulose, polyvinylpyrrolidone or gelatin. The liquid forms in suitably flavored suspending or dispersing agents may also include the synthetic and natural gums, for example, tragacanth, acacia, methyl-cellulose and the like. For parenteral administration, sterile suspensions and solutions are desired. Isotonic preparations which generally contain suitable preservatives are employed when intravenous administration is desired.

Advantageously, compounds of the present invention may be administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three or four times daily. Furthermore, compounds for the present invention can be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal skin patches well known to those of ordinary

skill in that art. To be administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen.

5

For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with an oral, non-toxic pharmaceutically acceptable inert carrier such as ethanol, glycerol, water and the like. Moreover, when desired or necessary, suitable binders; lubricants, disintegrating agents and coloring agents can also be incorporated into the mixture. Suitable binders include, without limitation, starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth or sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and the like. Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum and the like.

The daily dosage of the products may be varied over a wide range from 1 to 1000 mg per adult human per day. For oral administration, the compositions are preferably provided in the form of tablets containing, 0.01, 0.05, 0.1, 0.5, 1.0, 2.5, 5.0, 10.0, 15.0, 25.0, 50.0, 100, 150, 200, 250 and 500 milligrams of the active ingredient for the symptomatic adjustment of the dosage to the patient to be treated. An effective amount of the drug is ordinarily supplied at a dosage level of from about 0.01 mg/kg to about 30 mg/kg of body weight per day. Particularly, the range is from about 0.03 to about 15 mg/kg of body weight per day, and more particularly, from about 0.05 to about

10 mg/kg of body weight per day. The compounds may be administered on a regimen of 1 to 2 times per day.

Optimal dosages to be administered may be readily
 5 determined by those skilled in the art, and will vary
 with the particular compound used, the mode of
 administration, the strength of the preparation, the mode
 of administration, and the advancement of the disease
 condition. In addition, factors associated with the
 10 particular patient being treated, including patient age,
 weight, diet and time of administration, will result in
 the need to adjust dosages.

The compound of the present invention can also be
 15 administered in the form of liposome delivery systems,
 such as small unilamellar vesicles, large unilamellar
 vesicles, and multilamellar vesicles. Liposomes can be
 formed from a variety of lipids, including but not limited
 to amphipathic lipids such as phosphatidylcholines,
 20 sphingomyelins, phosphatidylethanolamines,
 phosphatidylcholines, cardiolipins, phosphatidylserines,
 phosphatidylglycerols, phosphatidic acids,
 phosphatidylinositols, diacyl trimethylammonium propanes,
 diacyl dimethylammonium propanes, and stearylamine,
 25 neutral lipids such as triglycerides, and combinations
 thereof. They may either contain cholesterol or may be
 cholesterol-free.

From Formula I it is evident that some of the
 30 compounds of the invention may have one or more
 asymmetric carbon atoms in their structure. It is
 intended that the present invention include within its
 scope the stereochemically pure isomeric forms of the
 compounds as well as their racemates. Stereochemically

pure isomeric forms may be obtained by the application of art known principles. Diastereoisomers may be separated by physical separation methods such as fractional crystallization and chromatographic techniques, and
5 enantiomers may be separated from each other by the selective crystallization of the diastereomeric salts with optically active acids or bases or by chiral chromatography. Pure stereoisomers may also be prepared synthetically from appropriate stereochemically pure
10 starting materials, or by using stereoselective reactions.

Some of the compounds of the present invention may have trans and cis isomers. In addition, where the
15 processes for the preparation of the compounds according to the invention give rise to mixture of stereoisomers, these isomers may be separated by conventional techniques such as preparative chromatography. The compounds may be prepared as a single stereoisomer or in racemic form as a
20 mixture of some possible stereoisomers. The non-racemic forms may be obtained by either synthesis or resolution. The compounds may, for example, be resolved into their component enantiomers by standard techniques, such as the formation of diastereomeric pairs by salt formation. The
25 compounds may also be resolved by covalent linkage to a chiral auxiliary, followed by chromatographic separation and/or crystallographic separation, and removal of the chiral auxiliary. Alternatively, the compounds may be resolved using chiral chromatography. The scope of the
30 present invention is intended to cover all such isomers or stereoisomers per se, as well as mixtures of cis and trans isomers, mixtures of diastereomers and racemic mixtures of enantiomers (optical isomers) as well.


During any of the processes for preparation of the compounds of the present invention, it may be necessary and/or desirable to protect sensitive or reactive groups on any of the molecules concerned. This may be achieved
 5 by means of conventional protecting groups, such as those described in Protective Groups in Organic Chemistry, ed. J.F.W. McOmie, Plenum Press, 1973; and T.W. Greene & P.G.M. Wuts, Protective Groups in Organic Synthesis, Third Edition, John Wiley & Sons, 1999. The protecting
 10 groups may be removed at a convenient subsequent stage using methods known from the art.

This invention will be better understood by reference to the schemes and examples that follow, but
 15 those skilled in the art will readily appreciate that these are only illustrative of the invention as described more fully in the claims which follow thereafter.


U.S. Pat. Nos. 5,696,117 and 5,854,242, both to
 20 Frechette et al., and WO 97/17333 (Frechette et al.) describe the synthesis of compounds of Formula I wherein E is hydrogen, all of which are hereby incorporated by reference.

25 The compounds of Formula I can be synthesized as outlined in the following schemes and equations.


Scheme 1



In accordance with Scheme 1, wherein Ar, Z^1 , Z^2 , A, G
 5 and n are as described hereinabove, the 4H-
 benzo[1,4]oxazin-3-ones can be made by conversion of the
 phenol of Formula 1, a known compound or a compound that
 may be prepared by known methods in the art, to the ether
 of Formula 2 and reduction to yield the corresponding
 10 compounds of Formula 3. Alternatively, compounds of
 Formula 3 can be obtained by reacting




with the aminophenol of Formula 4, a known compound or a compound that may be prepared by known methods in the art. Sequential protection of the primary alcohol and substitution of the amide produces the corresponding compounds of Formula 5. Deprotection of the alcohol gives the corresponding compounds of Formula 6 that can be further converted to the alkyl aryl ether of Formula Ia by adding HO-Ar-(CH₂)_nG' (G' is G as described above other than -COOH), which are commercially available and/or may be readily prepared by known methods. Thus deprotection of the ester to the acid yields the desired product of Formula Ib. The conditions for the above steps are more fully described in U.S. Pat. Nos. 5,696,117 and 5,854,242, both to Frechette et al., and WO 97/17333 (Frechette et al.).




To make compounds of Formula I wherein E is C₁-C₆ alkyl, compounds of Formula 5 can be exposed to a base such as NaH, LDA, or butyllithium followed by C₁-C₆ halide (Cl, Br, or I) to yield the corresponding compounds of 9 as shown in Equation 1 wherein P represents an appropriate protecting group. Compounds of 9 can be elaborated to final products as shown in Scheme 1. This procedure also applies to the compounds of Formula I wherein X is H.

When E is




compounds of Formula I can be obtained as shown in Equation 2 and Scheme 2, wherein P and P' are appropriate protecting groups and Z¹, Z², A, G and n are as described hereinabove, by replacement of α-bromo-γ-butyrolactone in
5 Scheme 1 with a reagent such as 10.



As represented by eq.2, treatment of γ -butyrolactone with a base, such as lithium diisopropylamide, and an electrophile, such as 2-(tert-butyldimethylsiloxy)ethyl bromide, then a base, such as lithium diisopropylamide, and a brominating agent, such as carbon tetrabromide, would yield the reagent **10**.

Scheme 2.




In accordance with Scheme 2, the 4H-benzo[1,4]oxanin-3-ones can be made by conversion of phenol **1** to ether **11** in the presence of the reagent **10** as described in Frechette, et al. The conversion can be carried out with a basic reagent such as an alkali metal carbonate or an alkali metal hydroxide, in a non-protic polar solvent, such as DMF (N,N-dimethylformamide) or THF, with or without heating. The resulting alkylaryl ether can be reduced to yield **12** with a reagent such as hydrogen gas or ammonium formate, and a catalyst, such as palladium or platinum, in an appropriate solvent, such as

methanol, ethanol, or ethyl acetate, at an appropriate temperature preferably between room temperature and 50 °C. Alternatively, compounds of Formula 12 can be obtained from aminophenol 4, by treatment of 4 with a base, such as an alkali metal hydride, in a non-protic polar solvent, such as DMF or THF, and the reagent 10, with or without heating. The primary alcohol can be protected with a reagent such as *tert*-butyldimethylsilyl chloride and imidazole in a non-protic polar solvent, such as DMF or THF, with or without heating.

Substitution of the amide by deprotonation with a base, such as an alkali metal hydride, in a non-protic polar solvent, such as DMF or THF, and addition of an alkyl halide can produce corresponding compounds of Formula 13. Selective deprotection of the alcohol can give compounds of Formula 14. The choice of deprotection methods will vary with the choice of protecting groups, using methods such as described in Protective Groups in Organic Chemistry, ed. J.F.W. McOmie, Plenum Press, 1973; and T.W. Greene & P.G.M. Wuts, Protective Groups in Organic Synthesis, Third Edition, John Wiley & Sons, 1999. The alkyl aryl ether 15 can be obtained by a reaction such as the Mitsunobu reaction as described by Frechette, et al. Deprotection of the alcohol and formation of the second alkyl aryl ether by a reaction such as the Mitsunobu reaction can yield compounds of Formula Ic. These procedures also apply to the compounds of Formula I wherein X is H.

Scheme 3



In accordance with Scheme 3, the 4H-benzo[1,4]oxanin-3-ones can be made by conversion of compounds of the Formula 17 (known or made) to compounds of the Formula 19 by methods in the literature. For example, the method of Buchwald (J. Am. Chem. Soc. 1996, 118, 7215) uses a palladium catalyst and compounds of the Formula 18. Cleavage of the methyl ether can be accomplished in a number of ways such as described in T.W. Greene & P.G.M. Wuts, Protective Groups in Organic Synthesis, Third Edition, John Wiley & Sons, 1999. The choice of deprotection methods may also be found in Protective Groups in Organic Chemistry, ed. J.F.W.

McOmie, Plenum Press, 1973; and T.W. Greene & P.G.M. Wuts, Protective Groups in Organic Synthesis, Third Edition, John Wiley & Sons, 1999. The resulting compounds of the Formula 20 can be utilized for the synthesis of the desired final product 22 with the method of U.S. Pat. Nos. 5,696,117 and 5,854,242, both to Frechette et al., and WO 97/17333 (Frechette et al.) outlined in Scheme 1. Thus, compounds of the Formula 20 can be treated with a base, such as an alkali metal hydride, in a non-protic polar solvent, such as DMF or THF, and 2-bromobutyrolactone, with or without heating, to provide compounds of the Formula 21. The compounds of the Formula Id can be obtained by a reaction such as the Mitsunobu reaction. These procedures also apply to the compounds of Formula I wherein X is H.

The following examples are intended to illustrate the invention but not to limit it.

Example 1


Intermediate 1: Dihydro-3-(2-nitrophenoxy)-2-(3H)-furanone

A mixture of 2-nitrophenol (20 g, 0.14 mol) and K₂CO₃ (25.2 g, 0.18 mol) in 280 mL DMF was cooled to 0 °C. 2-Bromobutyrolactone was added dropwise, the reaction was stirred for 45 min at 0 °C, then stirred at room temperature for 3 hours (h). The mixture was poured into 2 L water containing approx. 200 g salt, and the solution was washed with 6 X 100 mL of 1:1 diethyl ether/ethyl acetate. The combined organics were washed with 2 X 100 mL sat'd aq. K₂CO₃, 5 X 100 mL water and 100 mL brine, dried (Na₂SO₄), and filtered. Solvent was removed in

vacuo to yield the title compound as an off-white solid (21.88 g, 0.1 mol). ^1H (CDCl_3): 7.84 (d, 1H, $J = 7.9$), 7.57 (t, 1H, $J = 7.5$), 7.49 (d, 1H, $J = 7.9$), 7.16 (t, 1H, $J = 7.5$), 5.03 (t, 1H, $J = 7.4$), 4.58 (m, 1H), 4.42 (m, 1H), 2.8-2.6 (m, 2H).

Example 2


Intermediate 2: 2H-1,4-Benzoxazin-3(4H)-one, 2-[2-[(1,1-dimethylethyl)dimethylsilyl]oxy]ethyl]-



Intermediate 1 (21.88 g, 0.1 mol) was suspended in 200 mL ethanol and 200 mL EtOAc (ethyl acetate), then shaken overnight with 10% Pd/C and H_2 (45 psi) at room temperature. The solution was filtered through Celite and solvent was removed *in vacuo*. The crude product was dissolved in 200 anhydrous DMF, imidazole (14.1 g, 0.21 mol) was added, and the solution was cooled to 0 °C. *tert*-Butyldimethylsilyl chloride (31.2 g, 0.21 mol) was added as a solid and the reaction was stirred overnight, under N_2 , as the bath thawed. The reaction was poured into 1.4 L water containing approx. 200 g salt, and washed with 4 X 150 mL of 4:1 diethyl ether/EtOAc. The combined organics were washed with 6 X 100 mL water and 100 mL brine. The organics were dried (Na_2SO_4), filtered, and solvent was removed *in vacuo*. The product was isolated by silica gel chromatography with hexane/ethyl acetate. Obtain Intermediate 2 as a volatile solid (7.0 g, 0.022 mol). ^1H (CDCl_3): 6.89 (m, 3H), 6.80 (m, 1H), 4.70 (m, 1H), 3.78 (m, 2H), 2.15 (m, 1H), 1.94 (m, 1H), 0.83 (s, 9H), 0.07 (s, 6H).

Example 3


**Intermediate 3 (Method A): 2H-1,4-Benzoxazin-3(4H)-one,
2-[2-[[[(1,1-dimethylethyl)dimethylsilyl]oxy]ethyl]-
4-(5-oxohexyl)-**



A solution of Intermediate 2 (12.8 g, 0.04 mol) in 400 mL anh. DMF, under N₂, was cooled to 0 °C. Sodium hydride (75% dispersion in oil, 1.47 g, 0.046 mol) was added in two portions of 0.73 g with five minute intervals between additions. The solution was stirred for an additional 40 min at 0 °C. 6-Chloro-2-hexanone (5.6 g, 0.04 mol) in 20 mL DMF was added dropwise, the ice bath was replaced with an oil bath, and the solution was stirred at 65 °C for 17 h. The mixture was cooled to room temperature and poured into 1.5 L water containing approx. 200 g salt. The aqueous mixture was washed with 4 X 125 mL of 1:1 diethyl ether/ethyl acetate. The combined organics were washed with 6 X 125 mL water and 125 mL brine. The organics were dried (Na₂SO₄), filtered, and solvent was removed *in vacuo*. The product was isolated by silica gel chromatography with hexane/ethyl acetate. Intermediate 3 was obtained as a colorless oil (13.5 g, 0.034 mol). ¹H (CDCl₃): 6.92 (m, 4H), 4.63 (dd, 1H, *J* = 9.9, 3.7), 3.85-3.69 (m, 4H), 2.41 (br t, 2H), 2.17 (m and s, 4H), 1.93 (m, 1H), 1.58 (m, 4H), 0.82 (s, 9H), 0.07 (s, 6H).

Example 4


Intermediate 4 (Method B): 2H-1,4-Benzoxazin-3(4H)-one, 2-(2-hydroxyethyl)-4-(5-oxohexyl)-



Intermediate 3 was dissolved in 40 mL methanol and 4 mL water. Methanesulfonic acid (0.5 mL) was added and the mixture was stirred at room temperature for 2 h. Solvent was removed *in vacuo* and the product was isolated by silica gel chromatography with hexane/ethyl acetate. Intermediate 4 was obtained as a colorless oil (7.73 g, 0.027 mol). ^1H (CDCl_3): 6.77 (m, 4H), 4.51 (dd, 1H, $J = 8.9, 4.5$), 3.70 (m, 3H), 3.55 (m, 2H), 2.29 (br t, 2H), 1.91 (m and s, 4H), 1.78 (m, 1H), 1.42 (m, 4H).

Example 5

Compound 77 of Table 1 (Method C): Benzeneacetic acid, 2-[2-[3,4-dihydro-3-oxo-4-(5-oxohexyl)-2H-1,4-benzoxazin-2-yl]ethoxy]-, methyl ester



A solution of Intermediate 4 (8.75 g, 0.029 mol), methyl (2-hydroxyphenyl)acetate (7.48 g, 0.045 mol), and tributylphosphine (11.2 mL, 0.045 mol) in 500 mL anhydrous benzene, under N_2 , was cooled to 4 °C. 1,1'-(Azodicarbonyl)dipiperidine (11.34 g, 0.045 mol) was added in one portion, and the solution was stirred, with an overhead stirrer, at room temperature overnight. The

ORT-1527


organic phase was washed with 4 X 50 mL 2 N NaOH, 50 mL water and 50 mL brine. The organics were dried (Na₂SO₄), filtered, and solvent was removed *in vacuo*. The product

5 hexane/ethyl acetate. Compound 77 was obtained as a colorless oil (8.58 g, 0.02 mol). ¹H (CDCl₃): 7.28-6.89 (m, 8H), 4.76 (dd, 1H, *J* = 9.5, 4.0), 4.19 (m, 2H), 3.93 (br t, 2H), 3.61 and 3.60 (two singlets, 5H), 2.51 (m, 3H), 2.13 (m and s, 4H), 1.66 (m, 4H).

10

Example 6

Compound 78 of Table 1 (Method D): Benzeneacetic acid, 2-[2-[3,4-dihydro-3-oxo-4-(5-oxohexyl)-2H-1,4-benzoxazin-2-yl]ethoxy]-



15


A solution of Compound 77 (0.5 g, 1.13 mmol) in 10 mL THF was cooled to 0 °C. LiOH (0.143 g, 3.4 mmol) in 5 mL water was added in one portion. The solution was stirred at room temperature overnight, open to air. The solution was diluted with 25 mL water and 4 mL 1N HCl. Extract with dichloromethane. The combined organics were washed with water/brine combination, dried (Na₂SO₄), and filtered. Compound 78 was obtained as a colorless oil (0.41 g, 0.96 mmol). ¹H (CDCl₃): 7.28-6.89 (m, 8H), 4.83 (dd, 1H, *J* = 9.0, 3.6), 4.20 (m, 2H), 3.93 (m, 2H), 3.65 (d, 1H, *J* = 16.0), 3.59 (d, 1H, *J* = 16.0), 2.50 (m, 3H), 2.13 (m and s, 4H), 1.65 (m, 4H).

20

25

Example 7

Compound 79 of Table 1: Benzeneacetic acid, 2-[2-[3,4-dihydro-4-[(5Z)-5-(hydroxyimino)hexyl]-3-oxo-2H-1,4-benzoxazin-2-yl]ethoxy]-




Compound 78 (4.72 g, 0.011 mol) was dissolved in 200 mL ethanol and lutidine (2.6 mL, 0.022 mol).

Hydroxylamine hydrochloride (3.8 g, 0.055 mol) was added, and the reaction was stirred at room temperature for 2 hours. Solvent was removed, the residue was taken up in ethyl acetate and water, then further washed with 0.1 N HCl and water. The organics were dried (Na₂SO₄), filtered, and solvent was removed. Compound 79 was obtained as a white solid. ¹H (CDCl₃): 7.25 (m, 2H), 6.95 (m, 6H), 4.87 (dd, 1H, J = 9.5, 3.6), 4.27 (m, 2H), 4.15 (m, 1H), 3.91 (m, 1H), 3.65 (d, 1H, J = 15.3), 3.59 (d, 1H, J = 15.3), 2.50 (m, 1H), 2.4-2.1 (m, 3H), 1.9 (s, major isomer, approx. 3H) and 1.82 (s, minor isomer) 1.7-1.4 (m, 4H).

Example 8


Compound 81 of Table 1: Benzeneacetic acid, 2-[2-[3,4-dihydro-4-(5-hydroxyhexyl)-3-oxo-2H-1,4-benzoxazin-2-yl]ethoxy]-, methyl ester



A slurry of Compound 78 (0.2 g, 0.00046 mol) in 10 mL ethanol was treated with sodium borohydride (0.012 mg, 0.00031 mol), then stirred for 1 hour at room temperature. The reaction was diluted with water and extracted with ethyl acetate. The organics were dried (Na₂SO₄), filtered, and solvent was evaporated. Compound 81 was obtained as a colorless oil (0.052 g, 0.00012 mol). ¹H (CDCl₃): 7.25 (m, 2H), 6.95 (m, 6H), 4.76 (dd, 1H, *J* = 9.4, 3.9), 4.23 (m, 2H), 3.94 (t, 2H, *J* = 7.5), 3.80 (m, 1H), 3.61 and 3.60 (two singlets, 5H), 2.49 (m, 1H), 2.23 (m, 1H), 1.70 (m, 2H), 1.47 (m, 4H), 1.19 (d, 3H, *J* = 6.0).

Example 9

Compound 82 of Table 1: Benzeneacetic acid, 2-[2-[3,4-dihydro-4-(5-hydroxyhexyl)-3-oxo-2H-1,4-benzoxazin-2-yl]ethoxy]-




Prepared from Compound 81 (0.052 g, 0.00012 mol) in a manner analogous to Method D. Compound 82 was obtained as a colorless oil (0.040 g, 0.0001 mol). ¹H (CDCl₃): 7.25 (m, 2H), 6.95 (m, 6H), 4.87 (br d, 1H), 4.4-3.8 (m, 6H), 3.59 (s, 2H), 2.49 (m, 1H), 2.20 (m, 1H), 1.71 (m, 2H), 1.50 (m, 4H), 1.19 (m, 3H).

Example 10**Intermediate 5: 1-Bromo-6-fluorohexane**

(Diethylamino)sulfur trifluoride (4.0 mL, 0.031 mol) was cooled to 0 °C, and 6-bromohexan-1-ol (2.0 mL, 0.015 mol) was added dropwise. The mixture was heated to 35 °C for 4 h, then carefully poured into ice water. The aqueous phase was washed with methylene chloride. The organic phase was dried (Na₂SO₄), filtered, and solvent was removed *in vacuo*. 1-Bromo-6-fluorohexane was purified by silica gel chromatography with hexane/methylene chloride (2.1 g, 0.012 mol). ¹H (CDCl₃): 4.45 (dt, 2H, *J* = 47.3, 6.0), 3.42 (t, 2H, *J* = 6.6), 1.9-1.1 (m, 8H).

Example 11


Intermediate 6: 2H-1,4-Benzoxazin-3(4H)-one, 2-[2-[[[1,1-dimethylethyl)dimethylsilyl]oxy]ethyl]-4-(6-fluorohexyl)-



Prepared from Intermediate 2 (3.3 g, 0.011 mol) and 1-bromo-6-fluorohexane (2.0 g, 0.011 mol) in a manner analogous to Method A. Intermediate 6 was obtained as a colorless oil (2.9 g, 0.007 mol). MS (MH⁺): *m/z* = 410.

Example 12


Intermediate 7: 2H-1,4-Benzoxazin-3(4H)-one, 4-(6-fluorohexyl)-2-(2-hydroxyethyl)-



Prepared from Intermediate 6 (2.9 g, 0.007 mol) in a manner analogous to Method B. Intermediate 7 was obtained as a colorless oil (1.9 g, 0.006 mol). ^1H (CDCl₃): 7.02 (m, 4H), 4.69 (t, 1H, $J = 7.0$), 4.44 (dt, 2H, $J = 47.3, 6.0$), 3.88 (m, 4H), 2.37 (t, 1H, $J = 5.8$), 2.21 (m, 2H), 1.68 (m, 4H), 1.46 (m, 4H).

Example 13


Compound 90 of Table 1: Benzeneacetic acid, 2-[2-[4-(6-fluorohexyl)-3,4-dihydro-3-oxo-2H-1,4-benzoxazin-2-yl]ethoxy]-, methyl ester



Prepared from Intermediate 7 (1.8 g, 0.006 mol) in a manner analogous to Method C. Compound 90 was obtained as a colorless oil (2.2 g, 0.005 mol). ^1H (CDCl₃): 7.28-6.89 (m, 8H), 4.76 (dd, 1H, $J = 9.5, 4.0$), 4.44 (dt, 2H, $J = 47.3, 6.0$), 4.19 (m, 2H), 3.93 (br t, 2H), 3.61 and 3.60 (two singlets, 5H), 2.50 (m, 3H), 2.22 (m, 1H), 1.68 (m, 4H), 1.57 (m, 4H).

Example 14

Compound 91 of Table 1 (Method E): Benzeneacetic acid, 2-[2-[4-(6-fluorohexyl)-3,4-dihydro-3-oxo-2H-1,4-benzoxazin-2-yl]ethoxy]-



5

A solution of Compound 90 (2.1 g, 0.005 mol) in 25 mL methanol and 5 mL 2 N NaOH was heated to 55 °C. The solution was stirred for two hours, open to air. The solution was diluted with 25 mL water and 2 mL 6N HCl. Extract with dichloromethane. The combined organics were washed with water/brine combination, dried (Na₂SO₄), and filtered. Compound 91 was obtained as a colorless oil, then crystallized from pentane/diethyl ether (1.8 g, 0.004 mmol). ¹H (CDCl₃): 7.28-6.89 (m, 8H), 4.86 (dd, 1H, J = 9.1, 3.6), 4.20 (m, 2H), 4.44 (dt, 2H, J = 47.2, 6.1), 4.21 (m, 2H), 3.93 (br t, 2H), 3.66 (d, 1H, J = 16.0) and 3.59 (d, 1H, J = 16.0), 2.40 (m, 1H), 2.23 (m, 1H), 1.68 (m, 4H), 1.45 (m, 4H).


10

15

20

Example 15

Intermediate 8: 2H-1,4-Benzoxazin-3(4H)-one, 2-[2-[[[(1,1-dimethylethyl)dimethylsilyl]oxy]ethyl]-4-hexyl-



Prepared from Intermediate 1 (15.00 g, 0.049 mol) and 1-iodohexane (7.2 mL, 0.049 mol) in a manner analogous to Method A. Intermediate 8 was obtained as a colorless oil (18.03 g, 0.046 mol). ¹H (CDCl₃): 7.0 (m, 4H), 4.72 (dd,


25

ORT-1527

1H, $J = 10.0, 3.6$), 3.95-3.75 (m, 4H), 2.17 (m, 1H), 1.93 (m, 1H), 1.33 (m, 6H), 0.89 (s, 12H), 0.07 (s, 6H).

Example 16


5 **Intermediate 9: 2H-1,4-Benzoxazin-3(4H)-one,
4-hexyl-2-(2-hydroxyethyl)-**



Prepared from Intermediate 8 (18.3 g, 0.046 mol) in a manner analogous to Method B. Intermediate 9 was
10 obtained as a colorless oil (11.16 g, 0.04 mol). ^1H (CDCl₃): 7.01 (m, 4H), 4.69 (t, 1H, $J = 7.0$), 3.88 (m, 4H), 2.44 (t, 1H, $J = 5.8$), 2.20 (m, 2H), 1.65 (m, 2H), 1.33 (m, 6H), 0.89 (br t, 3H).

15 Example 17

Compound 14 of Table 1: Phosphonic acid, [[2-[2-(4-hexyl-3,4-dihydro-3-oxo-2H-1,4-benzoxazin-2-yl)ethoxy]phenyl]methyl]-, diethyl ester



20 Prepared from Intermediate 9 (1.13 g, 0.004 mol) and 2-(hydroxybenzyl)-phosphonic acid diethyl ester (*J. Org. Chem.* 1983, 48, 4768; 1.0 g, 0.04 mol) in a manner analogous to Method C. Compound 14 was obtained as a colorless oil (1.6 g, 0.003 mol). ^1H (CDCl₃): 7.4-6.7


ORT-1527

(m, 8H), 4.81 (dd, 1H, $J = 9.2, 4$), 4.4-3.9 (m, 8H), 3.24 (m, 2H), 2.55 (m, 1H), 2.27 (m, 1H), 1.64 (m, 2H), 1.5-1.2 (m, 14H), 0.89 (br t, 3H).

5

Example 18

Compound 15 of Table 1: Phosphonic acid, [[2-[2-(4-hexyl-3,4-dihydro-3-oxo-2H-1,4-benzoxazin-2-yl)ethoxy]phenyl]methyl]-



10

A solution of Compound 14 (0.5 g, 0.001 mol) in 25 mL 6N HCl was cooled to reflux overnight, open to air. Extract with 4 X 20 mL dichloromethane. The combined organics were washed with water and brine, dried (Na_2SO_4), and filtered. Solvent was removed *in vacuo*. Compound 15, contaminated with the half ester, was obtained as a colorless oil. (0.25 g, 0.0005 mol). ^1H (CDCl_3): 7.4-6.7 (m, 8H), 4.90 (m, 1H), 4.3-3.8 (m, 6H), 3.19 (m, 2H), 2.53 (m, 1H), 2.22 (m, 1H), 1.64 (m, 2H), 1.5-1.2 (m, 8H), 1.14 (t, from the half ester), 0.89 (br t, 3H). MS (M^-): $m/z = 474$.


15

20

Example 19

Intermediate 10: Benzeneacetic acid, 2-[2-(4-hexyl-3,4-dihydro-3-oxo-2H-1,4-benzoxazin-2-yl)ethoxy]-, methyl ester

25




Prepared from Intermediate 9 (11.16 g, 0.04 mol) in a manner analogous to Method C. Intermediate 10 was obtained as a colorless oil (9.7 g, 0.023 mol). ¹H

(CDCl₃): 7.28-6.89 (m, 8H), 4.76 (dd, 1H, *J* = 9.4, 4), 4.28-4.21 (m, 2H), 3.92 (t, 2H, *J* = 7.7), 3.60 (two singlets, 5H), 2.49 (m, 1H), 2.23 (m, 1H), 1.66 (m, 2H), 1.34 (m, 6H), 0.89 (br t, 3H).

Example 20

Intermediate 11: Benzeneacetic acid, 2-[2-(4-hexyl-3,4-dihydro-3-oxo-2H-1,4-benzoxazin-2-yl)ethoxy]-




Prepared from Intermediate 10 (26.5 g, 0.062 mol) in a manner analogous to Method D. Intermediate 11 was obtained by crystallization from pentane and diethyl ether (21.5 g, 0.052 mol). m.p. 80.0-81.5 °C. ¹H

(CDCl₃): 7.28-6.89 (m, 8H), 4.88 (dd, 1H, *J* = 9.0, 3.5), 4.20 (m, 2H), 3.91 (t, 2H, *J* = 7.8), 3.67 (d, 1H, *J* = 16.0), 3.60 (d, 1H, *J* = 16.0), 2.43 (m, 1H), 2.23 (m, 1H), 1.65 (m, 2H), 1.33 (m, 6H), 0.88 (br t, 3H).

Example 21


Compound 11 of Table 1 (Method F): Benzeneacetamide, 2-[2-(4-hexyl-3,4-dihydro-3-oxo-2H-1,4-benzoxazin-2-yl)ethoxy]-



Intermediate 11 (0.1 g, 0.00024 mol) was dissolved in 10 mL CH₂Cl₂ at room temperature. Carbonyl diimidazole (0.079 g, 0.0005 mol) was added and the stirred for 2 hours. Excess ammonia gas was added and the solution was stirred at room temperature for 30 minutes. The product was purified by silica gel chromatography with hexane/ethyl acetate. Compound 11 was obtained as a white solid (0.068 g, 0.00017 mol). ¹H (CDCl₃): 7.28-6.89, (m, 8H), 4.78 (dd, 1H, J = 8.2, 4.4), 4.25 (m, 2H), 3.91 (t, 2H, J = 7.1), 3.58 (d, 1H, J = 14.8), 3.52 (d, 1H, J = 14.8), 2.50 (m, 1H), 2.37 (m, 1H), 1.65 (m, 2H), 1.33 (m, 6H), 0.89 (br t, 3H).

Example 22

Compound 12 of Table 1: Benzeneacetic acid, 2-[2-(4-hexyl-3,4-dihydro-3-oxo-2H-1,4-benzoxazin-2-yl)ethoxy]-α-oxo-, methyl ester




Prepared from Intermediate 9 (0.61 g, 0.0022 mol) and 2-hydroxy-α-oxo-benzeneacetic acid, methyl ester (0.4

ORT-1527

g, 0.0022 mol). Compound 12 was obtained as a colorless oil (0.48 g, 0.0011 mol). $^1\text{H}(\text{CDCl}_3)$: 7.88 (d, 1H, $J = 7.4$), 7.58 (t, 1H, $J = 6.5$), 6.99 (m, 6H), 4.65 (dd, 1H, $J = 9.6, 3.9$), 4.57 (t, 2H, $J = 7.4$), 4.32 (m, 1H), 3.87 (s and m, 4H), 2.49 (m, 1H), 2.24 (m, 1H), 1.62 (m, 2H), 1.32 (m, 6H), 0.88 (br t, 3H).

Example 23

Compound 13 of Table 1: Benzeneacetic acid, 2-[2-(4-hexyl-3,4-dihydro-3-oxo-2H-1,4-benzoxazin-2-yl)ethoxy]-a-oxo-



Prepared from Compound 12 (0.2 g, 0.0005 mol) in a manner analogous to Method E. Compound 13 was obtained as a white solid from acetone (0.038 g, 0.0001 mol). $^1\text{H}(\text{CD}_3\text{OD})$: 7.83 (d, 1H, $J = 7.9$), 7.55 (t, 1H, $J = 6.9$), 7.02 (m, 6H), 5.00 (dd, 1H, $J = 9.7, 3.9$), 4.34 (m, 1H), 4.25 (m, 1H), 3.97 (m, 2H), 2.49 (m, 1H), 2.20 (m, 1H), 1.64 (m, 2H), 1.34 (m, 6H), 0.90 (br t, 3H).

Example 24

Intermediate 12: Benzyl 6-bromohexanoate

6-Bromohexanoic acid (10.0 g, 0.051 mol), benzyl alcohol (6.1 g, 0.056 mol) and DMAP (dimethylaminopyridine, 0.63 g, 0.005 mol) were dissolved in 100 mL CH_2Cl_2 , then cooled to 0 °C. 1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC, 10.8 g, 0.056 mol) was added and the reaction stirred overnight at room temperature. The organics were


ORT-1527

diluted with CH_2Cl_2 and washed with water, aq. NaHCO_3 , and brine. The organic phase was dried (Na_2SO_4), filtered, and solvent was removed. Intermediate 12 was obtained as a colorless oil (11.6 g, 0.04 mol). $\text{MS}(\text{MH}^+)$: $m/z = 286$.

5

Example 25

Intermediate 13: 2H-1,4-Benzoxazine-4-hexanoic acid, 2-[2-[[[(1,1-dimethylethyl)dimethylsilyl]oxy]ethyl]-3,4-dihydro-3-oxo-, phenylmethyl ester




10

Prepared from Intermediate 2 (7.5 g, 0.024 mol) and Intermediate 12 (7.0 g, 0.025 mol) in a manner analogous to Method A. Intermediate 13 was obtained as a colorless oil (13 g, 0.024 mol). $\text{MS}(\text{MH}^+)$: $m/z = 512$.

15

Example 26

Intermediate 14 (Method G): 2H-1,4-Benzoxazine-4-hexanoic acid, 3,4-dihydro-2-(2-hydroxyethyl)-3-oxo-, phenylmethyl ester



20


Intermediate 13 (13.7 g, 0.027 mol) was dissolved in 50 mL THF (tetrahydrofuran). Tetrabutylammonium fluoride

ORT-1527

(1.0 M in THF, 15.5 mL) was added and stirring continued under N₂ for 5 hours. Solvent was removed, the residue was taken up in ether and washed with water. The product was isolated by silica gel chromatography with hexane and ethyl acetate. Intermediate 14 was obtained as a colorless oil (10 g, 0.025 mmol). ¹H (CDCl₃): 7.34 (m, 5H), 7.00 (m, 4H), 5.10 (s, 2H), 4.68 (t, 1H, J = 5.6), 3.87 (m, 4H), 2.36 (t, 2H, J = 7.3), 2.19 (m, 2H), 1.67 (m, 4H), 1.37 (m, 2H).

Example 27


Compound 99 of Table 1: 2H-1,4-Benzoxazine-4-hexanoic acid, 3,4-dihydro-2-[2-[2-(2-methoxy-2-oxoethyl)phenoxy]ethyl]-3-oxo-, phenylmethyl ester



Prepared from Intermediate 14 (5.0 g, 0.013 mol) in a manner analogous to Method C. Compound 99 was obtained as a colorless oil (5.0 g, 0.009 mol). ¹H (CDCl₃): 7.4-6.8 (m, 13H), 5.11 (s, 2H), 4.75 (dd, 1H, J = 9.5, 3.9), 4.20 (m, 2H), 3.91 (t, 2H, J = 7.4), 3.61 and 3.59 (two singlets, 5H), 2.51 (m, 1H), 2.37 (t, 2H, J = 7.4), 2.22 (m, 1H), 1.73 (m, 4H), 1.41 (m, 2H).

Example 28


Compound 100 of Table 1: 2H-1,4-Benzoxazine-4-hexanoic acid, 3,4-dihydro-2-[2-[2-(2-methoxy-2-oxoethyl)phenoxy]ethyl]-3-oxo-



Compound 99 (5.0 g, 0.009 mol) was dissolved in 50 mL ethanol. The solution was exposed to 10% Pd/C and 50 psi H₂ for 4 hours. The solution was filtered through Celite and evaporated. Compound 100 was obtained as a colorless oil (3.7 g, 0.008 mol). ¹H (CDCl₃): 7.4-6.8 (m, 8H), 4.77 (dd, 1H, *J* = 9.5, 3.7), 4.18 (m, 2H), 3.93 (t, 2H, *J* = 7.1), 3.62 and 3.60 (two singlets, 5H), 2.48 (m, 1H), 2.34 (m, 2H), 2.22 (m, 1H), 1.68 (m, 4H), 1.43 (m, 2H).

Example 29


Compound 101 of Table 1: 2*H*-1,4-Benzoxazine-4-hexanoic acid, 2-[2-[2-(carboxymethyl)phenoxy]ethyl]-3,4-dihydro-3-oxo-



Prepared from Compound 100 (0.4 g, 0.0009 mol) in a manner analogous to Method E. Obtain Compound 101 as a white solid (0.25 g, 0.0006 mol). ¹H (CD₃OD): 7.3-6.8 (m, 8H), 4.83 (m, 1H), 4.19 (m, 2H), 3.97 (m, 2H), 3.58 (m, 2H), 2.5-2.1 (m, 4H), 1.64 (m, 4H), 1.40 (m, 2H).

Example 30


Compound 97 of Table 1: Benzenecetic acid, 2-[2-[4-(6-amino-6-oxohexyl)-3,4-dihydro-3-oxo-2H-1,4-benzoxazin-2-yl]ethoxy]-, methyl ester



Prepared from Compound 100 (0.4 g, 0.0009 mol) in a manner analogous to Method F. Obtain Compound 97 as a solid (0.3 g, 0.0007 mol). ^1H (CDCl_3): 7.4-6.8 (m, 8H), 4.76 (dd, 1H, $J = 9.3, 4.1$), 4.23 (m, 2H), 3.94 (t, 2H, $J = 7.2$), 3.62 and 3.59 (two singlets, 5H), 2.50 (m, 1H), 2.25 (m, 3H), 1.70 (m, 4H), 1.42 (m, 2H).

Example 31


Compound 98 of Table 1: Benzenecetic acid, 2-[2-[4-(6-amino-6-oxohexyl)-3,4-dihydro-3-oxo-2H-1,4-benzoxazin-2-yl]ethoxy]-



Prepared from Compound 97 (0.3 g, 0.0007 mol) in a manner analogous to Method D. Obtain Compound 98 as a white solid (0.24 g, 0.0005 mol). ^1H (DMSO): 7.3-6.8 (m, 8H), 4.79 (dd, 1H, $J = 9.2, 4.0$), 4.15 (m, 2H), 3.90 (t, 2H, $J = 7.2$), 3.50 (s, 2H), 2.27 (m, 1H), 2.02 (m, 3H), 1.50 (m, 4H), 1.28 (m, 2H).

Example 32

Intermediate 15: 2H-1,4-Benzoxazin-3(4H)-one, 2-[2-
 [[(1,1-dimethylethyl)dimethylsilyl]oxy]ethyl]-4-[[3-
 (trifluoromethyl)phenyl]methyl]-




Prepared from Intermediate 2 (1.38 g, 0.005 mol) and
 3-trifluoromethylbenzyl bromide (0.7 mL, 0.005 mol) in a
 manner analogous to Method A. Intermediate 15 was

obtained as a colorless oil (1.97 g, 0.004 mmol). ¹H
 (CDCl₃): 7.6-7.3 (m, 4H), 6.9-6.8 (m, 3H), 6.70 (d, 1H, J
 = 8 Hz), 5.10 (s, 2H), 4.79 (dd, 1H, J = 9.8, 3.7), 3.82
 (m, 2H), 2.20 (m, 1H), 1.93 (m, 1H), 0.82 (s, 9H), 0.0
 (s, 6H).

Example 33

Intermediate 16: 2H-1,4-Benzoxazin-3(4H)-one, 2-(2-
 hydroxyethyl)-4-[[3-(trifluoromethyl)phenyl]methyl]-




Prepared from Intermediate 15 (1.97 g, 0.004 mol) in
 a manner analogous to Method G. Intermediate 16 was

obtained as a colorless oil (1.14 g, 0.003 mmol). ¹H
 (CDCl₃): 7.6-7.3 (m, 4H), 6.9-6.8 (m, 3H), 6.70 (d, 1H, J
 = 8 Hz), 5.20 (s, 2H), 4.88 (t, 1H, J = 5.5), 3.82 (t,
 2H, J = 5.7), 2.4-2.1 (m, 3H).

Example 34


Compound 9 of Table 1: Benzoic acid, 2-[2-[3,4-dihydro-3-oxo-4-[[3-(trifluoromethyl)phenyl]methyl]-2H-1,4-benzoxazin-2-yl]ethoxy]-, methyl ester



Prepared from Intermediate 16 (0.183 g, 0.0005 mol) and methyl salicylate (0.119 g, 0.0008 mol) in a manner analogous to Method C. Compound 9 was obtained as a colorless oil (0.22 g, 0.0004 mmol). ^1H (CDCl_3): 7.81 (dd, 1H, $J = 7.7, 1.8$), 7.6-7.3 (m, 5H), 7.0-6.9 (m, 5H), 6.31 (d, 1H, $J = 7.4$ Hz), 5.21 (s, 2H), 5.08 (dd, 1H, $J = 9.5, 4.0$), 4.33 (m, 2H), 3.84 (s, 3H) 2.65 (m, 1H), 2.36 (m, 1H).

Example 35

Compound 26 of Table 1: Benzoic acid, 2-[2-[3,4-dihydro-3-oxo-4-[[3-(trifluoromethyl)phenyl]methyl]-2H-1,4-benzoxazin-2-yl]ethoxy]-, monosodium salt



Prepared from Compound 9 (0.182 g, 0.0004 mol) in a manner analogous to Method E, but after dilution of the completed reaction, product was isolated without addition of 6N HCl. The solid was filtered to yield Compound 26 as a sodium salt (0.141 g, 0.0003 mol). ^1H (DMSO): 7.81


ORT-1527

(dd, 1H, J = 7.7, 1.8), 7.6-7.3 (m, 5H), 7.0-6.9 (m, 5H), 6.81 (d, 1H, J = 7.4 Hz), 5.21 (s, 2H), 5.08 (dd, 1H, J = 9.5, 4.0), 4.33 (m, 2H), 3.84 (s, 3H) 2.65 (m, 1H), 2.36 (m, 1H).

5

Example 36

Compound 127 of Table 2: Benzeneacetic acid, 2-[2-(4-hexyl-3,4-dihydro-2-methyl-3-oxo-2H-1,4-benzoxazin-2-yl)ethoxy]-, methyl ester



10

Intermediate 11 (0.26 g, 0.0006 mol) was dissolved in 10 ml THF and cooled to 0 °C. Sodium hexamethyldisilazide (1.8 mL, 1.0 M) was added, then the reaction was stirred at room temperature for 30 minutes. The reaction was cooled to 0 °C, iodomethane (0.07 mL, 1.2 mmol) was added, and the reaction was stirred at room temperature overnight. The reaction was diluted with water and extracted with CH₂Cl₂. The organics were dried (Na₂SO₄), filtered, and solvent was removed in vacuo. The crude mixture, DMAP (0.004 g, 0.00003 mol) and methanol (0.05 mL, 0.001 mol) were dissolved in 20 mL CH₂Cl₂. 1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC, 0.059 g, 0.0003 mol) was added and the reaction stirred overnight at room temperature. Solvent was removed and the product was purified by silica gel chromatography. Compound 119 was obtained as a colorless oil (0.1 g, 0.00023 mol). ¹H (CDCl₃): 7.3-6.9 (m, 8H), 4.3-4.1 (m, 2H), 3.92 (m, 2H), 3.66 and 3.62 (two singlets, 5H), 2.49 (m, 1H), 2.25 (m, 1H), 1.66 (m, 2H), 1.51 (s, 3H), 1.33 (m, 6H), 0.89 (br t, 3H).


20

25

30

Example 37

Compound 128 of Table 2: Benzeneacetic acid, 2-[2-(4-hexyl-3,4-dihydro-2-methyl-3-oxo-2H-1,4-benzoxazin-2-yl)ethoxy]-




5

Prepared from Compound 127 (0.1 g, 0.00023 mol) in a manner analogous to Method D. Obtain Compound 120 as a colorless oil (0.078 g, 0.0002 mol). ^1H (CDCl_3): 7.3-6.8 (m, 8H), 4.22 (t, 2H, $J = 6.8$), 3.92 (m, 2H), 3.68 (d, 1H, $J = 15.5$), 3.59 (d, 1H, $J = 15.1$), 2.62 (m, 1H), 2.26 (m, 1H), 1.64 (m, 2H), 1.48 (s, 3H), 1.31 (m, 6H), 0.88 (br t, 3H).

10

Example 38

Compound 132 of Table 3: 2H-1,4-benzoxazin-3(4H)-one, 4-(4-methoxybutyl)-2-[2-[2-(1H-tetrazol-5-ylmethyl)phenoxy]ethyl]-, (2R)-



Compound 132


To a solution of the nitrile (370 mg, 0.94 mmol) in toluene (3 mL) was added sodium azide (80 mg, 1.22 mmol) and triethylamine hydrochloride (168 mg, 1.22 mmol). The reaction was heated at 100 °C for 20 h. The mixture was diluted with water and ethyl acetate (10 mL each) and acidified to pH=1 with conc. hydrochloric acid. The organic layer was washed with brine, dried (Na_2SO_4) and concentrated in vacuo to give 112 mg of the tetrazole as a white solid. MS: 460 ($\text{M}+\text{Na}$).

20

25


Example 39

Compound 126: Benzeneacetic acid, 2-[2-[3,4-dihydro-3-oxo-4-(6-methanesulfonamidoethyl)-2H-1,4-benzoxazin-2-yl]ethoxy]-



5

Intermediate 17: 2H-1,4-Benzoxazin-3(4H)-one, 2-[2-[[1,1-dimethylethyl]dimethylsilyloxy]ethyl]-4-(6-hexylphthalimide)-




A solution of Intermediate 2 (3.0 g, 9.8 mmol) in 20 mL anh. DMF, under N₂, was cooled to 0 °C. Sodium hydride (80% dispersion in oil, 0.35 g, 11.7 mmol) was added in two portions of 0.17 g with five minute intervals between additions. The solution was stirred for an additional 40 min at 0 °C. 6-bromo-hexylphthalimide (4.0 g, 12.7 mmol) in 5 mL DMF was added dropwise, the ice bath was replaced with an oil bath, and the solution was stirred at 50 °C for 17 h. The mixture was cooled to room temperature and poured into 1.5 L water containing approx. 200 g salt. The aqueous mixture was washed with 4 X 125 mL of 1:1 diethyl ether/ethyl acetate. The combined organics were

ORT-1527

washed with 6 X 125 mL water and 125 mL brine. The organics were dried (Na_2SO_4), filtered, and solvent was removed *in vacuo*. The product was isolated by silica gel chromatography with hexane/ethyl acetate. Intermediate 17 was obtained as a colorless oil (5.2 g, 9.8 mmol). MS: 559 (M+Na)

Intermediate 18: (2H-1,4-Benzoxazin-3(4H)-one, 2-(2-hydroxyethyl)-4-(6-hexylphthalimide)-




10

Intermediate 17 (6.2 g) was dissolved in 150 mL methanol and 20 mL water. Methanesulfonic acid (4 mL) was added and the mixture was stirred at room temperature for 2 h. Solvent was removed *in vacuo* and the product was isolated by silica gel chromatography with hexane/ethyl acetate. Intermediate 18 was obtained as a colorless oil (5.0 g, 11.4 mmol). MS: 445 (M+Na).

15

Intermediate 19: Benzeneacetic acid, 2-[2-[3,4-dihydro-3-oxo-4-(6-hexylphthalimide)-2H-1,4-benzoxazin-2-yl]ethoxy]-, methyl ester

20



25

A solution of Intermediate 18 (5 g, 11.8 mmol), methyl (2-hydroxyphenyl)acetate (3.9 g, 17.8 mmol), and tributylphosphine (4.4 mL, 17.8 mmol) in 180 mL anhydrous

ORT-1527

benzene, under N₂, was cooled to 4 °C. 1,1'-

(Azodicarbonyl)dipiperidine (4.5 g, 17.8 mmol) in THF (36 mL) was added in one portion, and the solution was

stirred, at room temperature overnight. The organic

5 phase was washed with 4 X 20 mL 2 N NaOH, 40 mL water and 40 mL brine. The organics were dried (Na₂SO₄), filtered,


and solvent was removed in vacuo. The product was

purified by silica gel chromatography with hexane/ethyl

acetate. Compound 77 was obtained as a yellow solid (2.6

10 g, 4.56 mmol). MS: 593 (M+Na)

Intermediate 20: Benzeneacetic acid, 2-[2-[3,4-dihydro-3-oxo-4-(6-methylsulfonamido)hexyl]-2H-1,4-benzoxazin-2-yl]ethoxy]-, methyl ester




To a suspension of intermediate 19 (1.0 g, 1.75 mmol) in 3.5 mL EtOH and 2 mL THF was added hydrazine (0.07 mL, 2.1 mmol). The reaction was stirred overnight
20 at room temperature, then diluted with MeOH and filtered. The solvent is evaporated in vacuo to give a white solid. To the white solid (0.2 g, 0.45 mmol) in 4 mL of CH₂Cl₂ is added methanesulfonyl chloride (0.04 mL, 0.6 mmol) and triethylamine (0.1 mL, 0.74 mmol). The reaction is
25 stirred for 17h at room temperature, then quenched with 2 mL of 1N HCl and diluted with 8 mL of EtOAc. The organic layer is washed with water and brine, dried (Na₂SO₄) and concentrated in vacuo. Purification by reverse-phase HPLC

ORT-1527

gave intermediate 20 as an orange oil (0.038g, 0.07 mmol). MS: 541 (M+Na).

5 Compound 126: Benzeneacetic acid, 2-[2-[3,4-dihydro-3-oxo-4-(6-methanesulfonamidohexyl)-2H-1,4-benzoxazin-2-yl]ethoxy]-



10 A solution of Intermediate 20 (0.038 g, 0.07 mmol) in 5 mL MeOH was cooled to 0 °C. NaOH (0.04 g, 1 mmol) in 1 mL water was added and the solution was stirred at room temperature overnight. The solution was acidified to pH=5 with 1N HCl. The mixture is extracted with ethyl acetate. The combined organics were washed with water/brine combination, dried (Na₂SO₄), and filtered.

15 Compound 126 was obtained as a yellow oil (0.03 g, 0.06 mmol). MS: 527 (M+Na).

20 Additional compounds in Tables 1, 2, and 3 below were made in the manners similar to the above examples and schemes.

aP2 Assay for Antagonist

25 Twenty-four hours after the initial seeding of the 96-well plates by hand (around 20,000/well), the differentiation assay may be initiated. Medium may be removed and replaced with 150µl of differentiation medium

containing vehicle (DMSO) or test compounds with a known aP2 activator or such aP2 activator alone. Cells may be returned to incubator for 24 hours culture. At the termination of the challenge, medium may be removed and 100 μ l of lysis buffer may be added to initiate the bDNA aP2 mRNA assay. The branched DNA assay may be performed according to the manufacturer's protocol (Bayer Diagnostics; Emeryville, CA). Result may be expressed as percent inhibition of aP2 mRNA production activated by the aP2 activator. IC_{50} 's may be determined by non-linear regression with a sigmoidal fit curve.

Following the challenge of the preadipocytes, cells may be lysed with lysis buffer (Bayer Diagnostics) containing the aP2 oligonucleotides. After a 15 minutes incubation at 53°C or 30 minutes at 37°C incubator, 70 μ l of the lysis buffer from each well may be added to a corresponding capture well (preincubated with 70 μ l of blocking buffer (Bayer Diagnostics)). The capture plate may be incubated overnight at 53°C in a plate incubator (Bayer Diagnostics). After this incubation, the bDNA and labeled probes may be annealed as directed by the manufacturer. Following a 30-minute incubation with the luminescent alkaline phosphatase substrate, dioxitane, the luminescence may be quantitated in a Dynex MLX microtiter plate luminometer. Oligonucleotide probes designed to anneal to the aP2 mRNA and function in the bDNA mRNA detection system are designed with *ProbeDesigner* software (Bayer Diagnostics). This software package analyzes a target sequence of interest with a series of algorithms in order to determine which regions of the sequence can perform as locations for


capture, label, or spacer probe annealing. The sequences of the oligonucleotides are as follows:

SEQ ID NO.1 CATTTTGTGAGTTTCTAGGATTATTCTTTTCTCTTGGAAGAAAGT
 SEQ ID NO.2 ATGTTAGGTTTGGCCATGCCTTTCTCTTGGAAGAAAGT
 SEQ ID NO.3 CCTCTCGTTTTCTCTTTATGGTTTTCTCTTGGAAGAAAGT
 SEQ ID NO.4 GCTTATGCTCTCTCATAAACTCTCGTGGTTTTCTCTTGGAAGAAAGT
 SEQ ID NO.5 CCAGGTACCTACAAAAGCATCACATTTAGGCATAGGACCCGTGTCT
 SEQ ID NO.6 GCCCACTCCTACTTCTTTTCATATAATCATTTAGGCATAGGACCCGTGTCT
 SEQ ID NO.7 AGCCACTTTCCTGGTGGCAAATTTAGGCATAGGACCCGTGTCT
 SEQ ID NO.8 CATCCCCATTACACTGATGATCTTTAGGCATAGGACCCGTGTCT
 SEQ ID NO.9 GTACCAGGACACCCCATCTAAGGTTTTTAGGCATAGGACCCGTGTCT
 SEQ ID NO.10 GGTTGATTTTCCATCCCATTTCTGCACATTTTAGGCATAGGACCCGTGTCT
 SEQ ID NO.11 GCATTCCACCACCAGTTTATCATTTTAGGCATAGGACCCGTGTCT
 SEQ ID NO.12 GCGAACTTCAGTCCAGGTCAACGTCCTTGTTTAGGCATAGGACCCGTGTCT
 SEQ ID NO.13 TCCACAGAATGTTGTAGAGTTCAATTTTAGGCATAGGACCCGTGTCT
 SEQ ID NO.14 AAAACAACAATATCTTTTGAACAATATAFTTAGGCATAGGACCCGTGTCT
 SEQ ID NO.15 TCAAAGTTTTCACTGGAGACAAGTTT
 SEQ ID NO.16 AAAGGTACTTTCAGATTTAATGGTGATCA
 SEQ ID NO.17 CTGGCCCAGTATGAAGGAAATCTCAGTATTTTT
 SEQ ID NO.18 TCTGCAGTGACTTCGTCAAATTC
 SEQ ID NO.19 ATGGTGCTCTTGACTTTCCTGTCA
 SEQ ID NO.20 AAGTGACGCCTTTCATGAC

5 aP2 Assay for Agonist

The procedure is described in detail in Burris et al., Molecular Endocrinology, 1999, 13:410, which is hereby incorporated by reference in its entirety, and aP2 assay results of agonist intrinsic activity may be presented as fold increase over vehicle in induction of aP2 mRNA production. Tables 1, 2, and 3 below set forth the mass spectra data and the agonist intrinsic activity of some compounds of the present invention.



Table 1

Compounds of this invention wherein Z¹ and Z² are both H

Compound No.	A	Pos(n)G	MS; MH ⁺ /Na	Agonist Intrinsic Activity
1	4-MeBn	2 (0) COOCH ₃	432	0.9
2	3-MeBn	2 (0) CF ₃	442	1.2
3	3-MeBn	2 (0) COOCH ₃	432	0.6
4	Bn	2 (0) COOCH ₃	418	1.0
5	4-ClBn	2 (0) COOCH ₃	452	0.7
6	3-ClBn	2 (0) COOCH ₃	452	-
7	3,5-Cl ₂ Bn	2 (0) COOCH ₃	486	-
8	3,4-Cl ₂ Bn	2 (0) CF ₃	497	3.3
9	3-CF ₃ Bn	2 (0) COOCH ₃	486	0.9
10	3-CF ₃ Bn	2 (0) CF ₃	496	0.8
11	n-Hex	2 (1) CONH ₂	411	-
12	n-Hex	2 (0) C(O) COOCH ₃	440	-
13	n-Hex	2 (0) C(O) COOH	426	2.6
14	n-Hex	2 (1) P(O) (OEt) ₂	503	1.6
15	n-Hex	2 (1) P(O) (OH) ₂	448	7.2
16	n-Hex	2 (0) CF ₃	422	-
17	n-Dec	2 (0) CF ₃	478	1.1
18	MeO (CH ₂) ₄	2 (0) CF ₃	424	0.3
19	EtO (CH ₂) ₃	2 (0) CF ₃	424	1.4
20	Me (CH ₂) ₂ O (CH ₂) ₂	2 (0) CF ₃	424	1.2
21	Me (CH ₂) ₂ S (CH ₂) ₂	2 (0) CF ₃	440	1.9
22	MeOC (O) CH ₂	2 (1) COOCH ₃	414	7.2
23	HOC (O) CH ₂	2 (1) COOH	386	0.8
24	AcOCH ₂ CH ₂	2 (1) COOCH ₃	428	1.3
25	HOCH ₂ CH ₂	2 (1) COOH	371	3.5
26	3-CF ₃ Bn	2 (0) COONa	472	3.1
27	3-CF ₃ Bn	2 (1) COOCH ₃	500	0.8
28	3-CF ₃ Bn	2 (1) COONa	486	21.5
29	3-CF ₃ Bn	2 (2) COOCH ₃	514	0.8
30	3-CF ₃ Bn	2 (2) COONa	500	5.6
31	3-CF ₃ Bn	2 (3) COOCH ₃	528	0.9
32	3-CF ₃ Bn	2 (3) COONa	514	8.2
33	3-CF ₃ Bn	3 (1) COOCH ₃	500	0.8
34	3-CF ₃ Bn	3 (1) COONa	486	10.4
35	3-CF ₃ Bn	3 (2) COOCH ₃	514	0.8
36	3-CF ₃ Bn	3 (2) COONa	500	11.1
37	4-CF ₃ Bn	2 (1) COOCH ₃	500	1.2
38	4-CF ₃ Bn	2 (1) COOH	486	38.7

39	3,5-(CF ₃) ₂ Bn	2(1) COOCH ₃	568	-
40	3,5-(CF ₃) ₂ Bn	2(1) COOH	554	6.7
41	3,5-(CF ₃) ₂ Bn	2(2) COOCH ₃	582	-
42	3,5-(CF ₃) ₂ Bn	2(2) COOH	568	2.1
43	3,5-(CF ₃) ₂ Bn	3(1) COOCH ₃	568	-
44	3,5-(CF ₃) ₂ Bn	3(1) COOH	554	6.8
45	MeOC(O)CH ₂	2(1) COOCH ₃	414	7.2
46	HOC(O)CH ₂	2(1) COOH	386	2
47	AcOCH ₂ CH ₂	2(1) COOCH ₃	428	1.3
48	HOCH ₂ CH ₂	2(1) COOH	371	8
49	propargyl	2(1) COOCH ₃	380	5.7
50	n-Hept	2(1) COOCH ₃	440	6.2
51	n-Hept	2(1) COOH	426	28.6
52	n-Oct	2(1) COOCH ₃	454	1.3
53	n-Oct	2(1) COOH	440	26.4
54	n-Non	2(1) COOCH ₃	468	0.9
55	n-Non	2(1) COOH	454	7.1
56	n-Dec	2(1) COOCH ₃	482	1.0
57	n-Dec	2(1) COOH	468	9.9
58	PhCH ₂ OC(O)(CH ₂) ₄	2(1) COOCH ₃	532	-
59	HOC(O)(CH ₂) ₄	2(1) COOCH ₃	442	-
60	HOC(O)(CH ₂) ₄	2(1) COOH	428	1.5
61	H ₂ NC(O)(CH ₂) ₄	2(1) COOCH ₃	441	-
62	H ₂ NC(O)(CH ₂) ₄	2(1) COOH	427	1.5
63	HO(CH ₂) ₅	2(1) COOCH ₃	428	-
64	HO(CH ₂) ₅	2(1) COOH	414	3.6
65	c-HexCH ₂ CH ₂	2(1) COOCH ₃	452	0.8
66	c-HexCH ₂ CH ₂	2(1) COOH	438	6.8
67	c-PentCH ₂ CH ₂	2(1) COOCH ₃	438	-
68	c-PentCH ₂ CH ₂	2(1) COOH	424	43.2
69	c-Pent(CH ₂) ₃	2(1) COOCH ₃	452	-
70	c-Pent(CH ₂) ₃	2(1) COOH	438	57.9
71	Me ₂ CH(CH ₂) ₄	2(1) COOCH ₃	440	-
72	Me ₂ CH(CH ₂) ₄	2(1) COOH	426	32
73	Me ₃ C(CH ₂) ₄	2(1) COOCH ₃	454	-
74	Me ₃ C(CH ₂) ₄	2(1) COOH	440	22.0
75	Me ₃ CCH ₂ CH(Me)(CH ₂) ₂	2(1) COOCH ₃	468	-
76	Me ₃ CCH ₂ CH(Me)(CH ₂) ₂	2(1) COOH	454	28.7
77	MeC(O)(CH ₂) ₄	2(1) COOCH ₃	440	9.9
78	MeC(O)(CH ₂) ₄	2(1) COOH	426	11.1
79	MeC(NOH)(CH ₂) ₄	2(1) COOH	441	4.5
80	MeC(NOMe)(CH ₂) ₄	2(1) COOH	455	2.8
81	MeCH(OH)(CH ₂) ₄	2(1) COOCH ₃	442	-
82	MeCH(OH)(CH ₂) ₄	2(1) COOH	428	4.9
83	Me ₂ C(OH)(CH ₂) ₄	2(1) COOCH ₃	456	-
84	Me ₂ C(OH)(CH ₂) ₄	2(1) COOH	442	3.5
85	MeCH(NH(Ac))(CH ₂) ₄	2(1) COOH	469	24
86	MeCF ₂ (CH ₂) ₄	2(1) COOCH ₃	462	-
87	MeCF ₂ (CH ₂) ₄	2(1) COOH	448	37.9

0990461.112101

88	enant 1; $\text{MeCF}_2(\text{CH}_2)_4$	2 (1) COOH	448	45.4
89	enant 2; $\text{MeCF}_2(\text{CH}_2)_4$	2 (1) COOH	448	42.0
90	$\text{F}(\text{CH}_2)_6$	2 (1) COOCH ₃	444	-
91	$\text{F}(\text{CH}_2)_6$	2 (1) COOH	430	64.9
92	enant 1; $\text{F}(\text{CH}_2)_6$	2 (1) COOH	430	44.8
93	enant 2; $\text{F}(\text{CH}_2)_6$	2 (1) COOH	430	53.1
94	$\text{HO}(\text{CH}_2)_6$	2 (1) COOCH ₃	442	-
95	$\text{HO}(\text{CH}_2)_6$	2 (1) COOH	428	36.9
96	$\text{H}_2\text{NCH}_2\text{CMe}_2(\text{CH}_2)_4$	2 (1) COOH	455	27.5
97	$\text{H}_2\text{NC}(\text{O})(\text{CH}_2)_5$	2 (1) COOCH ₃	455	-
98	$\text{H}_2\text{NC}(\text{O})(\text{CH}_2)_5$	2 (1) COOH	441	28
99	$\text{PhCH}_2\text{OC}(\text{O})(\text{CH}_2)_5$	2 (1) COOCH ₃	546	-
100	$\text{HOC}(\text{O})(\text{CH}_2)_5$	2 (1) COOCH ₃	456	-
101	$\text{HOC}(\text{O})(\text{CH}_2)_5$	2 (1) COOH	442	3.0
102	$\text{HOC}(\text{O})\text{CMe}_2(\text{CH}_2)_4$	2 (1) COOH	470	4.1
103	$\text{NCCMe}_2(\text{CH}_2)_4$	2 (1) COOCH ₃	465	-
104	$\text{NCCMe}_2(\text{CH}_2)_4$	2 (1) COOH	451	26.5
105	$\text{MeCF}_2(\text{CH}_2)_5$	2 (1) COOCH ₃	476	-
106	$\text{MeCF}_2(\text{CH}_2)_5$	2 (1) COOH	462	40.8
107	$\text{MeC}(\text{O})(\text{CH}_2)_5$	2 (1) COOCH ₃	454	-
108	$\text{MeC}(\text{O})(\text{CH}_2)_5$	2 (1) COOH	440	37.3
109	$\text{F}(\text{CH}_2)_7$	2 (1) COOCH ₃	458	-
110	$\text{F}(\text{CH}_2)_7$	2 (1) COOH	444	64.5
111	$\text{AcO}(\text{CH}_2)_7$	2 (1) COOCH ₃	498	-
112	$\text{HO}(\text{CH}_2)_7$	2 (1) COOH	442	74.5
113	$\text{F}(\text{CH}_2)_5$	2 (1) COOH	416	29.1
114	$\text{MeO}(\text{CH}_2)_4$	2 (1) COOH	414	50.7
115	$\text{EtO}(\text{CH}_2)_3$	2 (1) COOH	414	40.9
116	$\text{PrS}(\text{CH}_2)_2$	2 (1) COOH	430	4.0
117	$\text{PrO}(\text{CH}_2)_2$	2 (1) COOH	414	45.9
118	3,4-Cl ₂ Ph	2 (1) COOCH ₃	486	0.9
119	3,4-Cl ₂ Ph	2 (1) COOH	472	22.8
120		2 (1) COOH	513	22.5
121	n-Hex	4 (0) SC(CH ₃) ₂ COOH	472	16.13
122	$\text{CH}_3(\text{CH}_2)_2\text{NHCOCH}_2$	2 (1) COOH	449 (M+Na)	40.83
123	n-Hex	4 (0) 	480	1.65
124	$\text{CH}_3\text{NHCO}(\text{CH}_2)_3$	2 (1) COOH	449 (M+Na)	4.00
125	$\text{CH}_3\text{CONH}(\text{CH}_2)_6$	2 (1) COOH	491	10.78

ORT-1527



			(M+Na)	
126	$\text{CH}_3\text{SO}_2\text{NH}(\text{CH}_2)_6$	2 (1) COOH	527 (M+Na)	12.96

Keys:

Me=methyl; Et=ethyl; Pr=propyl; Bu=butyl; Pent=pentyl;
 Hex=hexyl; Hept=heptyl; Oct=octyl; Non=nonyl; Dec=decyl;
 5 Ac=acetyl; Bn=benzyl.

Table 2

Amides of this invention





Compound No.	Structure	MS; MH ⁺	Agonist Intrinsic Activity
127		440	-
128		426	2.5

0990461.112101

Table 3

Some stereoisomers of the invention

5

Compound No.	Structure	MS (MH ⁺ /M+Na)	PPAR γ Agonist Intrinsic Activity
129	 Chiral	456 (M+H)	12.62
130	 Chiral	448 (M+Na)	91.3
131	 Chiral	466 (M+Na)	36.4
132		460 (M+Na)	

03900461 44604

While the foregoing specification teaches the principles of the present invention, with examples provided for the purpose of illustration, it will be understood that the practice of the invention encompasses
5 all of the usual variations, adaptations and/or modifications as come within the scope of the following claims and their equivalents.

0930461-112101

ORT-1527

SEQUENCE LISTING

5 <110> Rybczynski, Philip
 et al.
 <120> BIOLOGICALLY ACTIVE 4H-BENZO[1,4]OXAZIN-3-ONES
 <130> pm431, patin2.1
 10 <140>
 <141>
 <160> 15
 15 <170> PatentIn Ver. 2.1
 <210> 1
 <211> 47
 <212> DNA
 20 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence: primer
 25 <400> 1
 cattttgtga gttttctagg attattcttt tctcttgga agaaagt 47
 <210> 2
 30 <211> 39
 <212> DNA
 <213> Artificial Sequence
 <220>
 35 <223> Description of Artificial Sequence: primer
 <400> 2
 atgttaggtt tggccatgcc tttctcttg aaagaaagt 39
 40 <210> 3
 <211> 41
 <212> DNA
 <213> Artificial Sequence
 45 <220>
 <223> Description of Artificial Sequence: primer
 <400> 3
 50 cctctcgttt tctctttatg gttttctctt ggaaagaaag t 41
 <210> 4
 <211> 47
 55 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> Description of Artificial Sequence: primer
 60 <400> 4

THE UNIVERSITY OF CHICAGO

47

5

10

46

15

20

25

30

35

43

40

45

45

50

55

<223> Description of Artificial Sequence: primer

60

48

09950451.112101

5 <210> 10
 <211> 51
 <212> DNA
 <213> Artificial Sequence

10 <220>
 <223> Description of Artificial Sequence: primer

<400> 10
 gggttgatttt ccatcccatt tctgcacatt ttaggcatag gacccgtgtc t 51

15 <210> 11
 <211> 45
 <212> DNA
 <213> Artificial Sequence

20 <220>
 <223> Description of Artificial Sequence: primer

<400> 11
 gcattccacc accagtttat catttttaggc ataggacccg tgtct 45

25 <210> 12
 <211> 52
 <212> DNA
 <213> Artificial Sequence

30 <220>
 <223> Description of Artificial Sequence: primer

35 <400> 12
 gcgaacttca gtccaggta acgtcccttg ttaggcata ggacccgtgt ct 52

40 <210> 13
 <211> 48
 <212> DNA
 <213> Artificial Sequence

45 <220>
 <223> Description of Artificial Sequence: primer

<400> 13
 tcccacagaa tggtgtagag ttcaatttta ggcataggac ccgtgtct 48

50 <210> 14
 <211> 51
 <212> DNA
 <213> Artificial Sequence

55 <220>
 <223> Description of Artificial Sequence: primer

60 <400> 14
 aaaacaacaa tatctttttg aacaatatat ttaggcatag gacccgtgtc t 51

ORT-1527

<210> 15

<211> 26

<212> DNA

5 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

10

<400> 15

tcaaagtttt cactggagac aagttt

26

09990461.1.13101